

LA-UR-21-23626

Approved for public release; distribution is unlimited.

Title: LANL Institutional Computing Annual Report Highlight: Local Time-Stepping in Global to Regional Ocean Modeling

Author(s): Petersen, Mark Roger
Capodaglio, Giacomo

Intended for: Report


Issued: 2021-04-14

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Local Time-Stepping in Global to Regional Ocean Modeling



LANL IC 2020 report, PI: Mark Petersen, Giacomo Capodaglio, CCS-2



Multi-resolution global ocean simulations are possible with the Model for Prediction Across Scales-Ocean (MPAS-Ocean), although the model is currently constrained to a uniform time step. The time step is determined by the size of the smallest cell in the grid, according to the CFL condition. To overcome this issue, a local time stepping (LTS) scheme has been designed for MPAS-Ocean. We have made progress on implementing the LTS scheme, as well as quantitative results and performance tests.

Local Time-Stepping in Global to Regional Ocean Modeling

LANL IC 2020 report, PI: Mark Petersen, Giacomo Capodaglio, CCS-2




The mesh is divided into regions with fine cells (blue), coarse cells (red) and interface layers (pink, yellow). The time steps subcycle on the fine region according to a positive number M , and conservatively communicate fluxes through the interface. If the coarse time-step is Δt , then the fine time-step is $\Delta t/M$. If $M=1$, no subcycling is done and the same time-step is used on all regions.


The algorithm has been implemented on global variable-resolution meshes with the shallow water equation set, which is a simple single-layer ocean model.

Local Time-Stepping in Global to Regional Ocean Modeling

LANL IC 2020 report, PI: Mark Petersen, Giacomo Capodaglio, CCS-2



Standard test cases with numerical reference solutions obtained with a Runge-Kutta scheme of order 4 are used to test accuracy and performance. Top plot shows convergence at third order, as expected, for the third order Local Time-Stepping scheme (LTS, yellow) with $M=1$, which matches up to machine precision the results obtained with the Strong Stability Preserving Runge-Kutta scheme on which it is based (red).



Performance curve comparison between a simulation with a uniform time step ($M=1$, blue and aqua) and a simulation where the fine region uses four sub-cycled time steps ($M=4$, orange and yellow). MB refers to a balancing only dictated by the cell sizes whereas LB assigns to each processor a balanced amount of fine, coarse and interface cells. Further improvement of the balancing procedure is under way.